

Cantaloupe Fungicide Spray Programs 2016

Bhabesh Dutta
Extension Vegetable Pathologist – University of Georgia

As with most crops, disease management begins prior to planting the crop to be protected. By using such methods as disease-free seed and transplants, proper rotation, disking and plowing, and use of resistant varieties, growers can minimize the amount of disease that is available to attack their crop. Using many practices takes all the weight off of any one practice and gives growers more disease management options at lower risk. Once plants are in the field, however, fungicide sprays are relied on heavily for effective and economical management of plant diseases. This spray guide is designed to encompass many diseases that attack cantaloupe during the growing season(s) in Georgia.

Spray Schedule = Start 1 week after transplanting, then use 7 day schedule if raining once a week.

Spring		Fall	
Spray No.	Fungicide(s)/disease activity	Spray No.	Fungicide(s)/disease activity
1.	Chlorothalonil/GSB copper at lowest labeled rate ¹ /BFB	1.	Chlorothalonil + tebuconazole ² /GSB copper at lowest labeled rate ¹ /BFB Presidio, Ranman, or Zampro/DOW
2.	Chlorothalonil/GSB copper at lowest labeled rate ¹ /BFB	2.	Chlorothalonil/GSB copper at lowest labeled rate ¹ /BFB
3.	Chlorothalonil/GSB copper at lowest labeled rate ¹ /BFB	3.	Orondis (soil app), Presidio, Revus/ PCAP
4.	Proline or Inspire Super or Aprovia top/GSB; Orondis (soil app), Presidio, Revus/ PCAP; Ranman or Previcur flex, or Curzate/DOW	3.	Chlorothalonil + tebuconazole ² OR Inspire Super or Aprovia top/GSB; copper at lowest labeled rate ¹ /BFB; Ranman or Previcur flex or Curzate/DOW
5.	Chlorothalonil + Luna Experience/GSB; Quintec or Torino or Proline or Procure/POW; Presidio or Revus / PCAP; Ranman or Previcur flex or Curzate/DOW	4.	Chlorothalonil/GSB Orondis (soil app) or Presidio, Revus/ PCAP; Ranman or Previcur flex or Curzate/DOW
6.	Proline or Inspire Super or Aprovia top/GSB; Quintec or Torino or Proline or Procure/POW; Orondis (soil app) or Presidio, Revus/ PCAP	5.	Chlorothalonil + tebuconazole OR Proline or Aprovia top/GSB; Quintec, Torino, Proline or Procure/POW; Presidio or Revus / PCAP; Ranman or Previcur flex, or Curzate/DOW
7.	Ranman or Previcur flex or Curzate/DOW Chlorothalonil + Luna Experience/GSB; Quintec, Torino, Proline or Procure/POW; Presidio or Revus/ PCAP; Ranman, Zampro, or Curzate/DOW	6.	Chlorothalonil + Luna experience/GSB; Quintec, Torino or Procure/POW; Presidio, Revus/ PCAP; Ranman or Previcur flex or Curzate/DOW
8 ³ .	Chlorothalonil/GSB; Quintec, Torino, Proline or Procure/POW; Presidio, Revus / PCAP; Ranman or Previcur flex or Curzate/DOW	7.	Chlorothalonil + tebuconazole OR Inspire Super/GSB Quintec, Torino or Procure/POW Presidio or Revus / PCAP Ranman, Zampro, or Curzate/DOW
		8 ³ .	Chlorothalonil + Fontelis/GSB + POW Quintec, Torino or Procure/POW Presidio, Revus / PCAP; Ranman or Previcur flex, or Curzate/DOW

¹ Copper is used prior to fruit set at lowest labeled rate if bacterial fruit blotch is an issue.

² Tebuconazole (any tebuconazole) should be used at 8.0 fl oz when sprayed each time with a seasonal limit of 24 fl oz.

³ Be mindful of preharvest intervals (PHIs) for later sprays as many may have a 7-14 day PHI.

Disease Acronyms: BFB = bacterial fruit blotch; GSB = gummy stem blight; POW = powdery mildew; DOW = downy mildew; ANTH = anthracnose; PCAP = Phytophthora crown and fruit rot.

Always read the label for detailed application instructions or consult the Georgia Pest Control Handbook

