

Supplemental Label



PRESIDIO[®] FUNGICIDE

EPA Reg. No. 59639-140
(Except California and New York)

PRESIDIO[®] FUNGICIDE FOR USE ON TOBACCO

This supplemental label expires on January 31, 2018 and must not be used or distributed after this date.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. READ THE LABEL AFFIXED TO THE CONTAINER FOR *PRESIDIO* FUNGICIDE BEFORE APPLYING. USE OF *PRESIDIO* FUNGICIDE ACCORDING TO THIS LABELING IS SUBJECT TO THE USE PRECAUTIONS AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR *PRESIDIO* FUNGICIDE.

| CROPS | ROTATIONAL INTERVAL |
|---|---------------------|
| Brassica (Cole) Leafy Vegetables Bulb Vegetables Cucurbit Vegetables Fruiting Vegetables Grapes Leafy Vegetables (Except Brassica) Root and Tuber Vegetables Tobacco | Immediately |
| Corn Wheat | 30 days |
| All Other Crops | 18 months |

SPECIFIC REQUIREMENTS

Restrictions and Limitations

- The maximum *Presidio* Fungicide application rate is 4 fl oz per acre per application and 12 fl oz per acre per season.
- Do not make more than 4 applications of *Presidio* Fungicide per acre per season unless otherwise noted under crop specific use instructions.

SPECIFIC REQUIREMENTS – continued

| Diseases | Application Rates | | Minimum Time from Last Application to Harvest (PHI) | Specific Use Instructions |
|--|----------------------|--------------------------|---|--|
| | fl oz/A | GPA Spray Mixture | | |
| Black Shank* (<i>Phytophthora nicotianae</i>) | 4 (0.125 lb ai/A) | Ground: Minimum 20 | N/A | <p>For resistance management, do not make sequential applications of <i>Presidio</i> Fungicide; alternate with a fungicide which has a different mode of action (different FRAC group) against the target pathogen.</p> <p>No more than 2 soil applications of <i>Presidio</i> Fungicide can be made per season.</p> <p>Minimum treatment interval 14 days.</p> <p><u>At-planting program</u> For control of black shank, apply preventively in transplant water (setter water) at planting. An additional application of <i>Presidio</i> Fungicide can be made as late as layby (last cultivation), if a fungicide with a different mode of action was used in the interim. When treating after transplanting, direct nozzles to cover soil beneath lower leaves; incorporate immediately with cultivator.</p> <p>or</p> <p><u>Post-transplant program</u> If a fungicide with a different mode of action is used at or immediately after transplanting, one application of <i>Presidio</i> Fungicide can be made at either first cultivation or layby (last cultivation). Direct nozzles to cover soil beneath lower leaves; incorporate immediately with cultivator.</p> |

*For best control of black shank with *Presidio* Fungicide, use with tobacco varieties that have moderate-to-high resistance to the black shank pathogen. Consult your local Cooperative Extension Service office or university specialist for information on variety selection.

SPECIFIC REQUIREMENTS – continued

| Diseases | Application Rates | | Minimum Time from Last Application to Harvest (PHI) | Specific Use Instructions |
|--|----------------------|--|---|--|
| | fl oz/A | GPA Spray Mixture | | |
| Blue Mold (<i>Peronospora tabacina</i>) | 4 (0.125 lb ai/A) | Ground minimum: 20 Aerial minimum: 10 | 7 days | For resistance management, <i>Presidio</i> Fungicide must be tank mixed with a labeled rate of another fungicide active against the target pathogen, but with a different mode of action. Apply as a foliar spray prior to disease onset or at first indication that blue mold is in the area. A second foliar application may be made, with a minimum treatment interval of 7 days between foliar applications. Make no more than 2 foliar applications per season. |

PLEASE CONTACT VALENT U.S.A. CORPORATION AT 800-6-VALENT (682-5368) TO DETERMINE IF THIS USE IS REGISTERED IN YOUR STATE.

©2015 Valent U.S.A. Corporation

Presidio is a registered trademark of Valent U.S.A. Corporation

Manufactured for:

Valent U.S.A. Corporation

P.O. Box 8025

Walnut Creek, CA 94596-8025

Made in U.S.A.

2015-PRD-0010

SAL20150112