

UGA Weed Control Programs for Broccoli, Cabbage, and Cauliflower 2015

A. S. Culpepper and J. C. Smith, University of Georgia, Tifton

Crop rotation, tillage, and a sound herbicide program are all often critical components for long-term success. This circular focuses on developing sound herbicide programs while minimizing crop injury for the following production systems: 1) transplanting into mulch, 2) transplanting into bareground, and 3) seeding into bareground. It is critical for growers to understand that their specific production practices may alter weed and crop responses; thus, **growers must evaluate these programs on limited acres until gaining experience.**

TRANSPLANT MULCH PRODUCTION: A systems approach using Goal 2XL, Dual Magnum, and Select Max can be extremely effective in the management of weeds while avoiding unacceptable crop injury.

Step 1. Control emerged weeds with Roundup and/or Gramoxone prior to planting; both products must be removed from mulch by 0.5 inch rain/irrigation in a single event prior to planting or severe injury can occur (Figure 1). Avoid Roundup applications within 7 d of planting even when washing mulch.

Step 2. Apply Goal 2XL prior to punching new holes and wait at least 5 days (**at least two days must be sunny**) after application **before** punching holes and transplanting. In most situations, Goal 2XL at 12 oz/A is in order. Goal is a very effective herbicide for controlling pigweeds, purslane, primrose, henbit, and many more broadleaf weeds (Figure 2).

Step 3. About 10 days after transplanting when plants are growing normally, apply Dual Magnum topically at 8 to 12 oz/A (Figure 3). Delay Dual Magnum application if plants are stressed.

Step 4. Apply Select Max at 8 to 12 oz/A without adjuvant when grasses are 3 inches or less.

Figure 1. Roundup must be removed from the mulch.



Roundup on mulch

No Roundup

Figure 2. Goal is very effective on many problem weeds.



Pigweed

No Goal

Goal preplant

TRANSPLANT BAREGROUND PRODUCTION: A systems approach using Treflan, Goal 2 XL, Dual Magnum, and Select Max can be extremely effective in the management of weeds while minimizing injury.


Step 1. Incorporate Treflan 2 inches deep into a moist soil at 10 to 16 oz/A; use lower rates on lighter soils, in an intense irrigation program, and/or in a harsh environment. Transplants are more tolerant than seeds.

Step 2. After land is prepared for planting, apply Goal 2XL, irrigate for ideal planting environment, wait at least 3 days (**at least two days must be sunny**), and then punch holes and transplant. Goal 2XL rate should be 8 to 12 oz/A; lower rates on sands with intense irrigation or when cold and wet. Also, use of transplants at least 5 weeks of age can reduce injury potential. Goal is a very effective herbicide for controlling pigweeds, purslane, primrose, henbit, and many more broadleaf weeds (Figure 2); include Gramoxone or Roundup when applying Goal 2XL if weeds have emerged.

Step 3. About 10 days after transplanting when plants are growing normally, apply Dual Magnum topically at a rate of 8 to 12 oz/A (Figure 3). Delay Dual Magnum application if plants are stressed.

Step 4. Apply Select Max at 8 to 12 oz/A without adjuvant when grasses are 3 inches or less.

Figure 3. Dual Magnum applied safely ovetop of cole crops 10 d after transplanting into mulch or bareground.



SEEDED BAREGROUND PRODUCTION: Seeded production offers greater challenges in weed control and poses greater crop injury potential as compared to transplants; however, the program below may be helpful.

Step 1. Incorporate Treflan 2 inches deep into a moist soil at 10 to 16 oz/A with lower rates used on lighter soils, in an intense irrigation program, and/or in a harsh environment.

Step 2. Once plants have reached 3 inches growing normally, apply Dual Magnum topically at 8 to 12 oz/A.

Step 3. Apply Select Max at 8 to 12 oz/A without adjuvant when grasses are 3 inches or less.

Critical Thinking Points To Avoid Crop Injury!

1. In transplant production, use the Goal 2XL formulation. **DO NOT USE GOAL IN SEEDED PRODUCTION!**
2. Use conservative herbicide rates when planting on sandy soil with low organic matter and/or with intense irrigation.
3. Dual Magnum topical application should be applied when crop is growing normally and not overly stressed.
4. In mulch, avoid Roundup over mulch within 7 d of planting regardless of washing the mulch. Gramoxone can be applied the day before planting as long as it is washed from the mulch appropriately.
5. In bareground production, do not apply Roundup within 3 days of transplanting.
6. In bareground transplant production, do not disturb Goal application with tillage.
7. Dual Magnum and Reflex are third party registrations and labels must be obtained from www.farmassist.com
8. Always follow label restrictions of each product used; read label for potential injury or carryover concerns.
9. Although growers prefer to spray Treflan preemergence after planting and then irrigate, this use pattern is not currently labeled. Efforts with Dow AgroSciences are underway to address this need.



THE UNIVERSITY OF GEORGIA
COOPERATIVE EXTENSION
 Colleges of Agricultural and Environmental Sciences & Family and Consumer Sciences

The University of Georgia and Ft. Valley State University, the U.S. Department of Agriculture and counties of the state cooperating. Cooperative Extension, the University of Georgia College of Agricultural and Environmental Sciences, offers educational programs, assistance and materials to all people without regard to race, color, national origin, age, gender or disability.

An Equal Opportunity Employer/Affirmative Action Organization
 Committed to a Diverse Work Force

Circular #

DATE

Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, The University of Georgia College of Agricultural and Environmental Sciences and the U.S. Department of Agriculture cooperating.
 J. Scott Angle, Dean and Director