
SUPPLEMENTAL LABELING

Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 1601A-S1A 0819

PYDIFLUMETOFEN	GROUP	7	FUNGICIDE
----------------	-------	----------	-----------

Miravis®

Fungicide

ADEPIDYN® Technology*

This supplemental label expires on 08/30/2022 and must not be used or distributed after this date.

Active Ingredient(s):	
Pydiflumetofen**	18.3%
Other Ingredients:	81.7%
Total:	100.0%

*Technology denotes the active ingredient Pydiflumetofen.

**CAS No. 1228284-64-7

Miravis is formulated as a suspension concentrate (SC) and contains 1.67 lb of pydiflumetofen per gallon.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

EPA Reg. No. 100-1601

All applicable directions, restrictions, and precautions on the EPA-registered label are to be followed. Before using Miravis as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA-registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and/or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.



DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

CROP USE DIRECTIONS Pome Fruit, Crop Group 11-10

Crops (Including all cultivars, varieties and/or hybrids of these)			
Apple	Mayhaw	Quince	
Azarole	Medlar	Quince, Chinese	
Crabapple	Pear	Quince, Japanese	
Loquat	Pear, Asian	Tejocote	
Target Disease	Rate fl oz/A (lb ai/A)	Application Timing	Use Directions
Alternaria blotch Alternaria rot <i>(Alternaria spp.)</i> Apple Scab <i>(Venturia inaequalis)</i> Cedar apple rust <i>(Gymnosporangium juniper-virginianae)</i> Flyspeck and Sooty blotch Pear Scab <i>(V. piris)</i> Powdery mildew <i>(Podosphaera leucotricha)</i> Quince rust <i>(Gymnosporangium spp.)</i>	3.4 (0.044)	<p><i>Protective Spray Schedule:</i> Apply every 7-10 days starting at ¼ to ½ inch green tip or when environmental conditions become conducive for scab. Continue through petal fall until the threat of primary scab is complete. For resistance management, combine Miravis with a protectant fungicide registered to control scab beginning at bloom.</p> <p><i>Curative Spray Schedule:</i> Use a forecasting system beginning at green tip. Apply within 48 hours of the onset of an infection period. Apply a follow up spray within 7 days. For resistance management, combine Miravis with a protectant fungicide registered to control scab beginning at bloom.</p> <p><i>Calendar Spray:</i> Apply the high rate of Miravis on a 14 day interval beginning at pink. Make no more than 2 consecutive applications before alternating to a non-Group 7 registered fungicide. Be sure to use according to that label.</p> <p>Rusts, leaf spots, summer diseases – Begin applications preventively. Apply Miravis alone or in combination with</p>	Apply by ground or air. An adjuvant may be added at recommended rates. Apply in sufficient volume to obtain coverage. If disease pressure is high, use the shortest interval.

		other non-Group 7 fungicides.	
Suppression Only: Bitter rot (<i>Glomerella cingulata</i>) Black rot (<i>Botryosphaeria obtusa</i>) Brooks fruit spot (<i>Mycosphaerella pomi</i>) White rot (<i>Botryosphaeria dothidea</i>)	3.4 (0.044)		
Resistance Management:			
<ul style="list-style-type: none"> • Refer to Section 3.2. • Do not make more than two consecutive applications of Miravis or other Group 7 fungicide before alternation with a fungicide that is not in Group 7. 			
USE RESTRICTIONS			
<ol style="list-style-type: none"> 1) Refer to Section 6.1 for additional product use restrictions. 2) Maximum Single Application Rate: DO NOT exceed the maximum rate listed in the table. 3) Minimum Application Interval: 7 days 4) Maximum Annual Rate: 13.6 fl oz/A/year <ol style="list-style-type: none"> a. DO NOT exceed 0.178 lb ai/A/year of pydiflumetofen-containing products. 5) DO NOT make more than 4 applications at the maximum application rate per year. 6) For aerial applications do not apply in less than 10 GPA water. 7) Pre-harvest Interval (PHI): 30 days 			

Stone Fruit, Crop Group 12-12

Crops (Including all cultivars, varieties and/or hybrids of these)			
Apricot	Nectarine	Plum, Chickasaw	
Apricot, Japanese	Peach	Plum, Damson	
Capulin	Plum	Plum, Japanese	
Cherry, black	Plum, American	Plum, Klamath	
Cherry, Nanking	Plum, beach	Plumcot	
Cherry, sweet	Plum, Canada	Plum, prune	
Cherry, tart	Plum, cherry	Sloe	
Jujube, Chinese			
Target Disease	Rate fl oz/A (lb ai/A)	Application Timing	Use Directions
Alternaria spot and fruit rot (<i>A. alternata</i>) Brown rot blossom blight and fruit rot (<i>Monilinia fructicola</i> , <i>M. laxa</i>) Powdery mildew (<i>Sphaerotheca pannosa</i> , <i>Podosphaera clandestina</i>) Scab (<i>Cladosporium carpophilum</i>) Shot hole (<i>Wilsonomyces carpophilus</i>)	5.1 (0.044 – 0.067)	For Brown rot blossom blight, begin applications at early bloom and continue through petal fall. For Brown rot on fruit, apply as needed a maximum of two sprays during the preharvest period up to the day of harvest (minimum of a 7-day retreatment interval). If high inoculum and severe disease conditions persist, apply a registered non-Group 7 fungicide. For all other diseases, follow the Brown rot blossom blight schedule. Make additional applications on a 7- to 14-day interval from the end of petal fall to harvest, following the resistance management guidelines.	Apply by ground or air. An adjuvant may be added at recommended rates. If disease pressure is high, use the shortest interval.
Resistance Management:			
<ul style="list-style-type: none"> Refer to Section 3.2. Do not make more than two consecutive applications of Miravis or other Group 7 fungicide before alternation with a fungicide that is not in Group 7. 			
USE RESTRICTIONS			
<ol style="list-style-type: none"> Refer to Section 6.1 for additional product use restrictions. Maximum Single Application Rate: DO NOT exceed the maximum rate listed in the table. Minimum Application Interval: 7 days Maximum Annual Rate: 20.4 fl oz/A/year <ol style="list-style-type: none"> DO NOT exceed 0.268 lb ai/A/year of pydiflumetofen-containing products. DO NOT make more than 4 applications at the maximum application rate per year. For aerial applications do not apply in less than 10 GPA water. Pre-harvest Interval (PHI): 0 days 			

Miravis®, Adepidyn™ and the Syngenta logo are trademarks of a Syngenta Group Company