




2019 Bayer Peanut Disease Risk Spray Schedules



Field Name: _____ Planting Date: _____



Low Risk		ABSOLUTE MAXX 3.5 oz -OR- Chlorothalonil 1.5 pt	28 Days	PROVOST SILVER 13 oz	28 Days	PROVOST SILVER 13 oz	28 Days	Chlorothalonil 1.5 pt						
Moderate Risk	PROLINE 5.7 oz In-Furrow	ABSOLUTE MAXX 3.5 oz -OR- Chlorothalonil 1.5 pt	21 Days	PROVOST SILVER 13 oz	21 Days	Non Group 3 White Mold Fungicide**	21 Days	PROVOST SILVER 13 oz	21 Days	Chlorothalonil 1.5 pt				
High Risk	PROLINE 5.7 oz In-Furrow	ABSOLUTE MAXX 3.5 oz -OR- Chlorothalonil 1.5 pt	14 Days	Chlorothalonil 1.5 pt	14 Days	Non Group 3 White Mold Fungicide**	14 Days	PROVOST SILVER 13 oz	14 Days	Non Group 3 White Mold Fungicide**	14 Days	PROVOST SILVER 13 oz	14 Days	Chlorothalonil 1.5 pt
CBR Program*	PROLINE 5.7 oz In-Furrow	Chlorothalonil 1.5 pt --OR-- PROLINE 5.7 oz Banded	30 Days	ABSOLUTE MAXX 3.5 oz -OR- Chlorothalonil 1.5 pt	14 Days	PROVOST SILVER 13 oz	14 Days	Non Group 3 White Mold Fungicide**	14 Days	PROVOST SILVER 13 oz	14 Days	Non Group 3 White Mold Fungicide**	14 Days	Chlorothalonil 1.5 pt
Nematode Program	VELUM TOTAL 18 oz In-Furrow	ABSOLUTE MAXX 3.5 oz -OR- Chlorothalonil 1.5 pt	45 Days	PROPULSE 13.6 oz	14 Days	PROVOST SILVER 13 oz	14 Days	Non Group 3 White Mold Fungicide**	14 Days	PROVOST SILVER 13 oz	14 Days	Chlorothalonil 1.5 pt		
Your Program														

See reverse side to assess your Peanut Disease Risk Index

Programs developed with the cooperation of:



* Fields with a history of or threat from *Cylindrocladium Black Rot* (CBR) should use the Bayer CBR disease management program coupled with a CBR resistant peanut variety.
 ** For resistance management, growers should rotate with non-DMI (Fungicide Group 3) fungicides. Do not use other DMI fungicides such as tebuconazole in these timings. If a grower chooses to use a strobilurin products such as pyraclostrobin or azoxystrobin in these timings, mix with other non-DMI fungicides such as chlorothalonil due to disease resistance. Contact your local Bayer rep for more information.

Under Peanut Rx, Bayer brand fungicides are the only fungicides that may be used in a grower program to qualify for Bayer standard product performance protection.

©2019 Bayer CropScience LP, 2 T.W. Alexander Drive, Research Triangle Park, NC 27709. Always read and follow label instructions. Bayer®, the Bayer Cross®, Proline®, Provost®, Absolute®, Velum Total® and Propulse® are registered trademarks of Bayer. The Peanut Rx logo and the UGA Extension logos are a trademark of The University of Georgia. The University of Florida IFAS logo is a trademark of the University of Florida. The Alabama Extension logo is a trademark of Auburn University. The Mississippi State University logo is a trademark of Mississippi State University. Not all products are currently registered for sale or use in all states. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our Web site at www.BayerCropScience.com.



Science For A Better Life



Assess Disease Risk in Your Field and Develop a Peanut Rx

This worksheet will lead you through the four-step process of determining your disease risk level in order to customize a Peanut Rx™ for your individual field. Use the reverse side of this worksheet with the assistance of your Bayer representative to develop a program specifically for your field.

For each of the risk index factors, identify which option best describes the situation in your field and add the index value associated with each choice to obtain your overall disease risk value. This worksheet does not contain all of the notes that accompany each factor included in the 2019 Peanut Rx. To view the complete 2019 Peanut Rx, visit the University of Georgia peanut web site at www.ugapeanuts.com.

Step 1: Assess Your Disease Risk

Variety Selection	TSWV	Leaf Spot	Soilborne Disease Points	
	Points	Points	White Mold	Limb Rot
AU NPL 17 ^{1,2}	15	15	15	NA
Bailey ³	10	25	10	NA
Florida Fancy ²	25	20	20	NA
FloRun™ '331' ²	10	20	15	NA
Georgia-06G	10	20	20	NA
Georgia-07W	10	20	15	NA
Georgia-09B ²	20	25	25	NA
Georgia-12Y ⁵	5	15	10	NA
Georgia-14N ^{2,4}	5	15	15	NA
Georgia-16HO ²	10	25	20	NA
Georgia Green	30	20	25	NA
Sullivan ^{1,2}	10	25	15	NA
Tiftguard ⁵	10	15	15	NA
TifNV-HiOL ^{2,4}	5	15	15	NA
TUFRunner™ '297' ²	10	25	20	NA
TUFRunner™ '511' ²	20	30	15	NA
Planting Date				
Prior to May 1	30	0	10	0
May 1 – May 10	15	5	5	0
May 11 – May 31	5	10	0	0
June 1 – June 10	10	15	0	5
After June 10	15	15	0	5
Plant Population (final stand, not seeding rate)				
Less than 3 plants per foot	25	NA	0	NA
3 to 4 plants per foot (for varieties with spotted wilt points greater than 25)	15	NA	0	NA
3 to 4 plants per foot (for varieties with spotted wilt points less than 25)	10	NA	0	NA
More than 4 plants per foot	5	NA	5	NA
At-Plant Insecticide				
None	15	NA	NA	NA
Other than Thimet 20G	15	NA	NA	NA
Thimet 20G	5	NA	NA	NA
Row Pattern				
Single rows	10	0	5	0
Twin rows	5	0	0	0
Tillage				
Conventional	15	10	0	0
Reduced	5	0	5	5
Classic Herbicide				
Classic applied	5	NA	NA	NA
No Classic applied	0	NA	NA	NA

¹Adequate research data is not available for all varieties with regards to all diseases. Additional varieties will be included as data to support the assignment of an index value are available.

²High oleic variety.

³Varieties Bailey have increased resistance to *Cylindrocladium black rot* (CBR) than do other varieties commonly planted in Georgia.

⁴Tiftguard and Georgia 14-N have excellent resistance to the peanut root-knot nematode.

⁵Georgia-12Y appears to have increased risk to *Rhizoctonia limb rot* and precautions should be taken to protect against this disease.

Crop Rotation with a Non-Legume Crop	TSWV	Leaf Spot	Soilborne Disease Points	
	Points	Points	White Mold	Limb Rot
0	NA	25	25	20
1	NA	15	20	15
2	NA	10	10	10
3 or more	NA	5	5	5
Field History				
No	NA	0	0	0
Yes	NA	10	15	10
Irrigation				
No	NA	0	0	0
Yes	NA	10	5	10

Step 2: Calculate Your Severity Points

Fill in following table to calculate your severity points for each of the four major peanut diseases given the 10 determining factors. Total each column to establish your disease index values.

Calculate Your Risk				
Add your index values for each determining factor below:	TSWV Points	Leaf Spot Points	White Mold Points	Limb Rot Points
Peanut Variety				
Planting Date				
Plant Population		----		----
At-Plant Insecticide		----	----	----
Row Pattern				
Tillage				
Classic Herbicide		----	----	----
Crop Rotation	----			
Field History	----			
Irrigation	----			
Your Total Index Value				

Step 3: Interpret Your Index Values

Once you've calculated your index values, utilize the table below to interpret your risk level.

Risk Index Category				
Risk Category:	TSWV Points	Leaf Spot Points	White Mold Points	Limb Rot Points
High Risk	≥ 115	65-100	55-80	TBD
Moderate Risk	70-110	40-60	30-50	TBD
Low Risk	≤65	10-35	10-25	TBD

In a year when tomato spotted wilt virus incidence is high statewide or in your region, even fields with a low risk level may experience significant losses. Consider the following recommendations to reduce your spotted wilt risk level: 1 - Use less susceptible varieties. 2 - Adjust your planting date. 3 - Consult the complete Peanut Rx for additional options that may provide limited benefit.

Step 4: Develop your Peanut Rx

Once you have calculated your total risk for each peanut disease, utilize the most conservative fungicide program as your guide for customizing a per field prescription spray program with the assistance of your Bayer CropScience representative. Bayer CropScience recommended disease risk spray schedules for each risk level are included on the reverse side of this worksheet.



Science For A Better Life