



UNIVERSITY OF GEORGIA
EXTENSION

2019 Commercial Pecan Spray Guide

Edited by Lenny Wells

Extension Pecan Specialist, Department of Horticulture

CONTRIBUTING AUTHORS

Tim Brenneman, Department of Plant Pathology

Jason Brock, Department of Plant Pathology

A. Stanley Culpepper, Extension Agronomist-Weed Science

Will Hudson, Extension Entomologist

Wayne Mitchem, Extension Associate-Weed Science

Lenny Wells, Extension Horticulturist

ONLINE RESOURCES

<http://blog.extension.uga.edu/pecan/>

www.caes.uga.edu/commodities/fruits/pecan/

<http://wiki.bugwood.org/Pecan/Georgia>

www.ent.uga.edu/pest-management/

It is important to always read any pesticide label before use. Use the product strictly according to the label directions. It is particularly important to follow all safety precautions. Trade and brand names are used only for information. The University of Georgia does not guarantee nor warrant published standards on any product mentioned; neither does the use of a trade or brand name imply approval of any product to the exclusion of others, which may also be suitable.

COMMERCIAL PECAN INSECT CONTROL (BEARING TREES)

Will Hudson, Extension Entomologist, and Angel Acebes, Research Entomologist

ORCHARD SURVEY PROCEDURES

Insect and mite infestation levels should be estimated at least weekly based on thorough orchard sampling. Sample trees in all segments of each orchard. A good method is to sample every fourth tree in every fourth tree row (about 10% of the trees). Sample each major cultivar represented in the orchard. Sample a minimum of 10 terminals per tree. Check all compound leaves and the nut clusters on each terminal. Check as high in the tree as possible. Foliar pest

counts should be made on compound leaves surrounding the nut clusters. Nut clusters should be inspected carefully for the presence of pests or damage. Hickory shuckworm damage should be monitored mid-season by examining fallen nuts for a whitish spot on the side. Pecan weevil populations should be monitored by survey traps.

PEST	PESTICIDE	MOA	AMOUNT PER ACRE	REI/PHI (Hours or Days)	TIMING AND REMARKS
Phylloxera	<i>chlorpyrifos</i> 4E Lorsban, Chlorphos	1B	2 pt	24 H/ –	Treat trees with a recent history of heavy infestation and surrounding trees. Apply at budbreak with the first pre-pollination spray. Note: Other <i>imidacloprid</i> formulations are available. Read labels carefully to find the proper rate.
	Centric 40WG	4A	2-2.5 oz	12 H/ –	
	<i>imidacloprid</i> Provado 1.6F Trimax Pro	4A	3.5 oz 1.3-2.6 oz	12 H/ –	
Spittlebugs	<i>imidacloprid</i> Trimax, Provado, many generics	4A	See label <i>Several formulations are available.</i>	12 H/ –	Spittlebug infestations are easily recognized by the white, frothy masses on terminals or nut clusters. Definite thresholds have not been established and treatment is seldom needed. Many generic <i>imidacloprid</i> formulations are available.
Pecan Nut Casebearer	<i>chlorpyrifos</i> 4E Lorsban, Chlorphos	1B	1.5 pt	24 H/ –	Light infestations causing occasional damage do not require control in most crop years. The most serious damage usually occurs in mid-May. Adult emergence should be monitored with pheromone traps. Place traps in orchards by mid-April. Begin sampling for nut casebearer in the first week of May. Pay particular attention to orchards not under a spray program the preceding year and orchards with a recent history of nut casebearer problems. Try to time sprays to stop injury before more than one nut per cluster is infested. It is recommended that broad-spectrum contact insecticides, such as <i>chlorpyrifos</i> and the pyrethroids, not be used in early- or mid-season to conserve beneficial insect populations. (See Special Considerations section.)
	<i>methoxyfenozide</i> Intrepid 2F	18	4-8 oz	4 H/ –	
	<i>spinosad</i> Spintor 2SC	5	4-10 oz	4 H/ –	
	<i>diflubenzuron</i> Dimilin 2L	15	8-16 oz	12 H/ –	
	<i>clothianadin</i> Belay	4A	3-6 oz	12 H/ –	
	<i>methoxyfenozide</i> + <i>spinetoram</i> Intrepid Edge	5 + 18	4-6.4 oz	4 H –	
	<i>tolfenpyrad</i> Apta	21	17-27 oz	12 H –	DO NOT apply more than 1 application. No more than 27 oz/A/season.
<i>abamectin</i> + <i>cyantraniliprole</i> Minecto Pro	6 + 28	8-12 oz	12 H/ 21 D	No more than 2 consecutive applications, no more than 24 oz/season.	

COMMERCIAL PECAN INSECT CONTROL

PEST	PESTICIDE	MOA	AMOUNT PER ACRE	REI/PHI (Hours or Days)	TIMING AND REMARKS
Mites	<i>abamectin</i> Agri-Mek SC and others	6	2.25-4.25 oz	12 H/ -	A non-ionic surfactant or horticultural oil MUST be added to the tank.
	<i>bifenazate</i> Acramite 4SC	Unclassified	12-16 oz	12 H/ -	Mites, especially the pecan leaf scorch mite, are normally late season pests. Mite damage appears as bronzed, scorched areas on the undersides of leaflets. Scorched areas begin at the leaflet midribs then spread out toward leaflet margins. Mites often build up on low limbs in the shaded, interior portions of trees then spread rapidly up and out. For heavy infestations, repeat the application in 5-7 days.
	<i>spirodiclofen</i> Envidor 2SC	23	14-18 oz	12 H/ -	
	<i>fenpyroximate</i> Portal	21A	2 pt	12 H/ -	
	<i>pyridaben</i> Nexter	21	5.2-10.67 oz	24 H/ -	
	<i>hexythiazox</i> Savey 50DF	10A	3-6 oz	12 H/ -	
	<i>etoxazole</i> Zeal	10B	2-3 oz	12 H/ -	
	<i>fenazaquin</i> Magister SC	21	24-36 oz	12 H/ 7 D	No more than one application per year.
Yellow Aphids	FOLIAR APPLICATIONS				
	<i>acetamiprid</i> Assail 30SG	4A	2.5-9.6 oz	12 H/ -	Yellow aphids may be present in orchards throughout the growing season. Populations are usually highest in April-May and again in August-September. In early season, DO NOT treat yellow aphids if they are the only insect problem. Rely on beneficial insects to suppress early season populations.
	<i>clothianidin</i> Belay	4A	3-6 fl oz	12 H/ -	
	<i>flonicamid</i> Beleaf, Carbine	9C	2-2.8 oz	12 H/ -	
	<i>flupyradifurone</i> Sivanto 200 SL	4D	7.0-10.5 oz	4 H/ 7 D	In prolonged dry periods, lower, chronic aphid populations may require treatment to prevent the build-up of unacceptable levels of honeydew and sooty mold. WEEKLY SCOUTING IS VERY IMPORTANT IN TIMING APHID SPRAYS, ESPECIALLY IN LATE SEASON. Rotate among classes of insecticides between treatments to avoid resistance development.
	<i>imidacloprid</i> Provado, many generics	4A	See label	12 H/ -	
	<i>pymetrozine</i> Fulfill	9B	4 oz	12 H/ -	It is suggested that pyrethroid materials (<i>cypermethrin</i> , <i>bifenthrin</i> , etc.) not be used, alone or in combination, in early- or mid-season applications.
	<i>pyridaben</i> Nexter	21	5.2-10.67 oz	24 H/ -	
	<i>sulfoxaflor</i> Closer	4C	1.5-2.75 oz	12 H/ 7 D	Many generic formulations of <i>imidacloprid</i> are available. Read label carefully for recommended rate. <i>Imidacloprid</i> alone may not control yellow and black-margined aphids.
	<i>thiamethoxam</i> Centric	4A	2-2.5 oz	12 H/ -	
	<i>tolfenpyrad</i> Apta	21A	17-27 oz	12 H/ -	
	SYSTEMIC APPLICATIONS				
<i>imidacloprid</i> Admire Pro	4A	7-14 fl oz	12 H/ -	Admire can be applied through a drip irrigation system, as an emitter spot application, or as a shanked-in emitter adjacent application. See label for complete details. Apply Admire only to orchards where drip irrigation has been established for at least 5 years. DO NOT apply more than 1 application of Apta, no more than 27 oz/A/season.	
					Use the 14 oz rate for black pecan aphid control.

COMMERCIAL PECAN INSECT CONTROL

PEST	PESTICIDE	MOA	AMOUNT PER ACRE	REI/PHI (Hours or Days)	TIMING AND REMARKS
Black Pecan Aphid	SAME INSECTICIDES AS FOR YELLOW APHIDS or <i>chlorpyrifos</i> Lorsban, generics	1B	Check label	24 H/ –	Black pecan aphids may cause damage as early as May but are usually a serious problem only in late season. Damage appears as yellow spots on leaflets. Damaged spots later turn brown and 2-4 damaged spots per leaflet can cause leaflet drop. Carefully check all compound leaves on 10 terminals per tree, on at least 10 trees per orchard for the presence of black pecan aphids. Prior to July 1, treat if 25% of terminals have 2 or more black aphids. After July 1, treat if 15% of terminals have more than one black aphid and nymph clusters are found. Concentrate checks on susceptible cultivars such as Schley, Sumner, and Gloria Grande. Be sure to check all compound leaves on each terminal examined.
	<i>gibberellic acid</i> ProGibb 4% ProGibb LV Plus	N/A	10 oz 5 fl oz	N/A	<i>Gibberellic acid</i> is a plant growth regulator that prevents damage from black pecan aphid feeding and inhibits establishment in the orchard. It does not affect aphids directly and will not control any other pest, including yellow aphids. Three applications should be made at 2-week intervals, beginning in mid-July, applying 10 oz (or 5 oz of ProGibb LV Plus) each time.
Hickory Shuckworm	<i>chlorpyrifos</i> 4E Lorsban, Chlorfos	1B	1-14 pt	24 H/ –	Shuckworms are active throughout the season, but do not cause significant damage until June or later. Prior to shell hardening, larval feeding causes nuts to drop. After shells harden, feeding causes shucks to stick to the shells, reducing quality. If orchards have a history of shuckworm infestation, a spray should be applied in early June. In early August, 2-3 additional sprays should be applied. Initiate August sprays at half-shell hardening and repeat at 2-week intervals until shuck split if shuckworm activity continues. <i>Chlorpyrifos</i> and pyrethroids (Asana, Ambush, Mustang, etc.) applied for other pests will also control shuckworm. It is not necessary to spray in August if pecan weevil controls are applied. Please note the Special Considerations section regarding the use of pyrethroid materials. DO NOT apply more than 1 application, no more than 27 oz/A/season.
	<i>clothianadin</i> Belay	4A	3-6 oz	12 H/ –	
	<i>diflubenzuron</i> Dimilin 2L	15	8-16 oz	12 H/ –	
	<i>methoxyfenozide</i> Intrepid 2F, Turnstyle	18	4-8 oz	4 H/ –	
	<i>methoxyfenozide</i> + <i>spinetoram</i> Intrepid Edge	5 + 18	4-6.4 oz	4 H/ –	
	<i>tolfenpyrad</i> Apta	21A	17-27 oz	12 H/ –	
	<i>abamectin</i> + <i>cyantraniliprole</i> Minecto Pro	6 + 28	8-12 oz	24 H/ 21 D	
	<i>chlorantraniliprole</i> + <i>lambda-cyhalothrin</i> Besiege	3 + 28	6-12.5 oz	24 H/ –	Besiege contains a pyrethroid, and may flare aphids and mites if used in early or mid-season. The best fit is for late season shuckworm.

COMMERCIAL PECAN INSECT CONTROL

PEST	PESTICIDE	MOA	AMOUNT PER ACRE	REI/PHI (Hours or Days)	TIMING AND REMARKS
Pecan Weevil	<i>carbaryl</i> Carbaryl 80S Sevin	1A	3 lb	24 H/ –	<p>Pecan weevil emergence may extend from July into October. Peak emergence is normally between August 10 and September 20. Emergence should be monitored in each infested grove with traps, knockdown sprays or a combination of these methods. Trees known to have a recent history of weevil problems should be selected for monitoring. If excessive nut drop results from pecan weevil feeding punctures before pecan shells begin to harden, spray at once. After pecan shells harden and nuts reach the “dough” or “gel” stage, treat when weevils emerge (especially following rains) and continue at 7-10 day intervals until emergence stops. APHID OR MITE POPULATIONS MAY BUILD UP WHERE <i>CARBARYL</i> IS USED. If these pests become a problem, apply aphicides or miticides as previously directed.</p> <p>NOTE: Several pyrethroids, (Asana, Ammo, Baythroid, Brigade, Mustang Max) as well as <i>Imidan</i> are labeled for pecan weevil control. If these materials are used for weevils, they can be expected to be most effective where weevil populations are low. They may be adequate to prevent feeding injury from weevils emerging prior to shell hardening but their use could be risky under heavy weevil pressure after nuts reach the gel stage and are subject to weevil oviposition. (See Special Considerations section).</p> <p>Several products are available that combine a pyrethroid insecticide with an aphicide. These products may help suppress aphids while providing weevil control. Brand names include Endigo, Leverage, and others.</p>
	<i>carbaryl</i> Carbaryl 4F Sevin XLR Various pyrethroids	1A	4-5 qt	24 H/ –	

COMMERCIAL PECAN INSECT CONTROL

KERNAL FEEDING HEMIPTERANS

(Stink bugs and Plant bugs)

A complex of true bugs (stink bugs and plant bugs) attack pecan. They may be present in orchards all year but normally cause their most serious injury from late August through September. Prior to shell hardening, feeding injury causes nut drop. After shell hardening, their feeding causes black, bitter spots on kernels, reducing quality. They can continue to feed, through the hardened shells, until nuts are harvested. The presence and numbers of stink bugs and plant bugs should be noted in surveys throughout the season. Special attention should be paid to the true bugs in late-season orchard surveys. Treat when 1 stink bug is found per 40 terminals OR when 5 or more are found per knockdown spray on a sheet covering 20% of the area under a tree. Sprays for these insects are difficult to time properly because the bugs move in and out of orchards. Close checking is required to detect damaging populations. No materials have consistently given excellent stink bug control, possibly due to the difficulty in timing sprays. The pyrethroids are labeled for stink bug control. Please note the pre-harvest use restrictions of the products.

FIRE ANTS

Fire ants have been known to protect pecan aphids by destroying beneficial insects in pecan orchards. Fire ants should be controlled or at least kept out of pecan trees. Lorsban 4E at 2 pts/A as a ground spray is labeled for fire ant control. Best approach is probably applying an ant bait in late spring.

SCALE INSECTS

Scale populations build slowly, but can reach damaging levels before becoming obvious. Examine fallen limbs carefully during the season for scale presence. Preferred treatment is 1-2% horticultural oil spray, applied in November-December and again in February. For severe problems, an application of Esteem in June may be necessary.

OTHER INSECT PESTS

Pests such as pecan leaf casebearer, leaf miners, walnut caterpillar, fall webworm, pecan budmoth, nut curculio, shoot curculio, Prionus root borers, and others may occasionally cause economic injury to pecan. Growers should be able to identify these pests and their damage. Color photographs of all pecan pests and their injury can be found in the Southern Pecan Growers Handbook and online from the UGA Extension pecan team (Google search “ugapecans”). The publication is available at \$30 per copy. For ordering information, visit: extension.uga.edu/publications/for-sale.html

Specific controls for occasional pests not covered in this spray guide can be obtained from your local county Extension agent.

SPECIAL CONSIDERATIONS

Alternative Formulations – Some pesticides listed in this publication are available in formulations other than the ones listed. If different formulations are used, apply an equivalent amount of actual toxicant per acre.

Pest Resistance and Chemical Use – The aphids and mites which attack pecan have demonstrated the ability to become resistant to insecticides applied for their control. The rate at which this resistance develops depends on the chemical used, the frequency of use, the duration of use, and the rates used. Aphid and mite exposure to effective materials should be minimized to prolong the effective life of the chemicals. It is suggested that no insecticide be applied until it is absolutely necessary (this can be determined by thorough sampling) and that chemicals be alternated as much as possible. Resistance to *neonicotinyl* insecticides has developed in some areas for both yellow- and black-margined pecan aphids. This class of insecticides includes *imidacloprid*, *thiamethoxam*, *acetamiprid*, and *clothianidin*. These materials no longer provide adequate control of resistant populations. Aphid and mite populations may flare following application of Sevin or pyrethroids. Growers should be alert for this response, and limit applications of these materials to the minimum necessary for weevil or stink bug control.

Supplemental Control Measures – Beneficial insects such as lady beetles and lacewings provide natural assistance in suppressing aphid and mite populations. Beneficials are of particular value in early season. Elimination of unneeded early-season insecticide sprays conserves existing populations of beneficial insects and reduces the potential for severe aphid problems later in the season. The planting of leguminous cover crops in tree-row middles promotes the build-up and retention of lady beetle populations in orchards. Crimson clover and Hairy vetch appear to be two of the best ground covers. If leguminous ground covers are planted, an herbicide strip should be maintained down each tree row and special attention should be paid to the increased water requirements that are likely to exist. Extraneous plant material resulting from the heavy growth of legumes must be removed or broken down prior to harvest or implementation of a program of row middle vegetation suppression (see Weed Control section).

COMMERCIAL PECAN INSECT AND DISEASE SPRAY GUIDE

(NON-BEARING TREES)

Will Hudson, Extension Entomology
Jason Brock and Tim Brenneman, Plant Pathology

FOLIAR SPRAYS

TIME OF APPLICATION	PEST	PESTICIDE	MOA	AMOUNT PER ACRE	REI/PHI (Hours or Days)	INSTRUCTIONS AND REMARKS
Bud Break When first buds open.	Foliar disease	Fungicide + <i>chlorpyrifos</i> Chlorphos, Lorsban	1B	+ half rate 1-2 pt 4-8 oz	24 H/ -	Spray sufficient gallonage for thorough coverage. For fungicide options, refer to the Pre-pollination section for Pecan Disease Control.
	Pecan bud moth	<i>methoxyfenozide</i> Intrepid 2F	18	3-4 oz	4 H/ -	The phosphorous acid fungicides are particularly useful with their excellent activity on foliar diseases, highly systemic nature, and low risk of fungicide resistance.
		<i>methoxyfenozide</i> + <i>spinetoram</i> Intrepid Edge	5 + 18	4-6.4 oz	4 H/ -	
		<i>abamectin</i> + <i>cyantraniliprole</i> Minecto Pro	6 + 28	8-12 oz	12 H/ -	
	Hickory shoot curculio	<i>chlorpyrifos</i> Lorsban, Chlorphos, etc.	1B	1.5-2 pt	24 H/ -	Apply sprays for shoot curculio at bud-break on the earliest cultivars and repeat at 10-14 day intervals.
Cover Sprays Three weeks after bud-break spray and every 4-6 weeks as needed.	Foliar disease	Fungicide + <i>chlorpyrifos</i> Chlorphos, Lorsban	1B	See above + 1-2 pt	24 H/ -	Spray sufficient gallonage for thorough coverage.
	Pecan bud moth	<i>chlorpyrifos</i> Chlorphos, Lorsban, etc.	1B	1.5-2 pt	24 H/ -	
		<i>diflubenzuron</i> Dimilin 2L		8-16 oz	24 H/ -	
		Imidan 70WSP		1.5 lb		
		<i>methoxyfenozide</i> Intrepid 2F	18	4-8 oz	4 H/ -	
		<i>abamectin</i> + <i>cyantraniliprole</i> Minecto Pro	6 + 28	8-12 oz	12 H/ -	

PECAN CHEMICALS: PRE-HARVEST INTERVALS AND OTHER RESTRICTIONS

CHEMICAL	MOA	REI/PHI (Hours or Days)	TIMING AND REMARKS
Acramite 4 SC	Undetermined	12 H/ 14 D	Only 1 spray per year.
Admire	4A	12 H/ –	Apply to soil between May 15 and July 15. Apply only to orchards that have been established on trickle irrigation for at least 5 years. DO NOT apply more than 32 fl oz of Admire per acre per season as a soil application. DO NOT apply more than 0.5 lb ai of Admire or Provado/A/season.
Ammo		–/ 21 D	Up to 0.8 lb ai/A/season may be applied prior to shuck split. DO NOT graze or feed cover crops.
Asana		–/ 21 D	DO NOT feed or graze livestock on treated orchard floors. DO NOT exceed 0.3 lb ai/A/season. DO NOT mix with fungicides containing <i>triphenyltin hydroxide</i> .
Assail	4A	12 H/ 14 D	DO NOT apply more than 4 times per season, nor more often than every 7 days.
Baythroid		–/ 14 D	No more than 2.8 fl oz/A/season.
Belay	4A	12 H/ 21 D	No more than 12 oz/season. DO NOT graze.
Carbaryl	1A	24 H/ 14 D	DO NOT apply more than a total of 15 qt/season.
Centric	4A	12 H/ 14 D	DO NOT exceed 5 oz/A/season. Allow at least 7 days between applications.
Closer		–/ 7 D	No more than 4 applications per season, and no more than 2 consecutive applications.
Desperado		–/ 7 D	No more than 2.2 gal/season; no aerial application.
Dimethoate		–/ 21 D	DO NOT graze livestock in treated groves.
Elast F			DO NOT apply after shucks open. DO NOT graze treated areas.
Enable		–/ 28 D	DO NOT apply after shuck split. DO NOT apply more than 48 oz/A. DO NOT graze treated areas.
Endosulfan			DO NOT apply after shuck split. DO NOT graze livestock in treated groves. DO NOT exceed 2 applications per year or 4 qt/A/year.
Envidor	23	12 H/ 7 D	Maximum of 1 application per season.
Fury/Mustang		–/ 21 D	DO NOT apply more than 0.3 lb ai/A/season or after shuck split. DO NOT graze or cut treated cover crops for feed.
Headline		–/ 14 D	DO NOT apply more than 28 fl oz/A/season.

PECAN CHEMICALS: PRE-HARVEST INTERVALS AND OTHER RESTRICTIONS

CHEMICAL	MOA	REI/PHI (Hours or Days)	TIMING AND REMARKS
Imidan		3 D/ 14 D	DO NOT graze livestock in treated groves.
Intrepid	18	18/ 14 D	DO NOT graze livestock in treated areas or feed cover crops grown in treated areas. DO NOT apply more than 10 fl oz/application or 64 oz/season.
Kelthane		-/ 7 D	Applicators must be in enclosed cabs or cockpits.
Lorsban, Chlorphos	1B	24 H/ 28 D	DO NOT allow livestock to graze in treated orchards. Make no more than 5 applications per season.
Nexter	21A	24 H/ 7 D	No more than 10.67 oz/application nor more than 2 applications per season. No aerial applications.
Portal	21A	12 H/ 14 D	No more than one application per season.
Propimax			DO NOT apply after shuck split. DO NOT graze livestock in treated areas or cut treated areas for feed. DO NOT apply more than 32 fl oz/A/season.
Provado	4A	12 H/ -	DO NOT apply more than 28 fl oz of Provado/A/year. DO NOT apply more than a total of 0.5 lb ai of Provado or Admire/A/season.
Quilt		-/ 45 D	DO NOT apply after shuck split. DO NOT graze livestock in treated areas or cut treated areas for feed. DO NOT apply more than 122 fl oz/A/season.
Savey	10A	12 H/ -	DO NOT graze livestock in treated areas. Only one application per season may be made.
Sovran		-/ 45 D	DO NOT apply more than 25.6 fl oz/A/season.
Stratego		-/ 30 D	DO NOT apply after shuck split. DO NOT apply more than 30 fl oz/A/season.
Sulfur			No time limitations.
TPTH			DO NOT use more than 45 oz (36 oz ai) of product per season. DO NOT apply after shucks begin to open. DO NOT graze dairy or meat animals in treated groves.
Topsin M			DO NOT apply after shuck split. DO NOT graze livestock in treated areas or cut treated areas for feed. DO NOT apply more than 3 lb/A/season.
Trimax Pro	4A	12 H/ 7 D	Maximum of 10.1 oz/A allowed per crop season. Allow at least 10 days between applications.
Zeal	10B	12 H/ 28 D	Maximum of 1 application per season.

****DO NOT** graze livestock in treated groves where prohibited or until grazing restrictions have been met.

PECAN DISEASE CONTROL

Jason Brock and Tim Brenneman, Department of Plant Pathology

DISEASE	CHEMICAL & FORMULATION	MOA	RATE/ACRE	REI/PHI (Hours or Days)	COMMENTS
PRE-POLLINATION APPLICATIONS: Every 10-14 Days From Bud Break Through Nut Set					
Scab; Downy Spot	<i>azoxystrobin</i> Abound Azaka	11	6-12 fl oz	4 H/ 45 D	See MOA info on next page.
	<i>difenoconazole + azoxystrobin</i> Quadris Top	3 + 11	10-14 fl oz	12 H/ 45 D	
	<i>difenoconazole + azoxystrobin</i>	3 + 11	8-14 fl oz	12 H/ 21 D	
	<i>dodine</i> Elast 400F + FRAC Group 3 fungicide	U12 + 3	25 fl oz + half rate	48 H/ Do not apply after shuck split	For any tank mix combination of Elast, TPTH, or a Group 3 fungicide, the rates provided are the lowest recommended and will provide excellent control of scab under most conditions. When disease pressure is elevated, the rate of either mixing partner can be increased. See MOA info on next page.
	<i>dodine</i> Elast 400F + TPTH	U12 + 30	25 fl oz + half rate	48 H/ Do not apply after shuck split or within 30 D of harvest	For any tank mix combination of Elast, TPTH, or a Group 3 fungicide, the rates provided are the lowest recommended and will provide excellent control of scab under most conditions. When disease pressure is elevated, the rate of either mixing partner can be increased. See MOA info on next page.
	<i>fenbuconazole</i> Enable 2F	3	8 fl oz	12 H/ Do not apply after shuck split or within 28 D of harvest	See MOA info on next page.
	<i>kresoxim-methyl</i> Sovran	11	2.4-3.2 fl oz	12 H/ 45 D	
	<i>metconazole</i> Quash	3	3.5 oz	12 H/ 25 D	
	<i>phosphorous acid</i> <i>Kphite 7LP</i> Phostrol ProPhyt FungiPhite Reliant	33	2-6 pt 2-5 pt 2-3 pt 2-3 pt 4 pt	4 H/ -	
	<i>phosphorous acid + tebuconazole</i> Viathon	33 + 3	2-2.5 pt	12 H/ 0 D	
	<i>propiconazole</i> Orbit Propimax EC Bumper 41.8EC	3	8 fl oz	12 H/ Do not apply after shuck split	
	<i>propiconazole + azoxystrobin</i> Quilt Quilt Xcel	3 + 11	14-27.5 fl oz 14-21 fl oz	12 H/ Do not apply after shuck split or within 45 D of harvest	

PECAN DISEASE CONTROL

DISEASE	CHEMICAL & FORMULATION	MOA	RATE/ACRE	REI/PHI (Hours or Days)	COMMENTS
PRE-POLLINATION APPLICATIONS: Every 10-14 Days From Bud Break Through Nut Set					
Scab; Downy Spot <i>(continued)</i>	<i>pyraclostrobin</i> Headline	11	6-7 fl oz	12 H/ 14 D	<p>MOA Group 1: Risk for resistance is high. Use should be limited. When conditions are very favorable for scab, Topsin can be used in combination with either a full rate of TPTH or Elast. Limit the use to 1 or 2 applications per season.</p> <p>MOA Group 3: Resistance risk is moderate. For best results, tank mix <i>tebuconazole</i> with a surfactant. Do not add a surfactant if mixing with other fungicides. Increasing the rate of a Group 3 fungicide will be important if reduced sensitivity is known or suspected. Stand-alone use is not recommended where reduced sensitivity is known or suspected.</p> <p>MOA Group 11: Resistance risk is moderate. Do not make more than 2 sequential applications. If only using solo products, Group 11 fungicides should not be used in more than 1/3 of the total number of fungicide applications. If using Group 3 tank-mixed with other modes of action, they should not be used in more than 1/2 of the total number of fungicide applications.</p> <p>MOA Group 30: Resistance risk is low.</p> <p>MOA Group 33: Resistance risk is low. For best control apply in 100 gpa by ground. Do not apply in consecutive applications. Three to five applications are generally recommended. Check labels for potential limitations on maximum number of applications or amount of active ingredient allowed per season. Do not use when there is a <i>phosphate</i> deficiency.</p> <p>MOA Group U12: Resistance risk is low. Do not use on Moore, Van Deman, Barton, or Shawnee. Do not use a surfactant. Do not use with foliar zinc treatments.</p>
	<i>tebuconazole</i> Folicur 3.6F Tebuzole 3.6F Monsoon Orius 3.6F Toledo 3.6F	3	8 fl oz	12 H/ Do not apply after shuck split	
	<i>tetraconazole</i> Andiamo	3	8.5 fl oz	12 H/ 30 D	
	<i>tetraconazole + azoxystrobin</i> Brixen	3 + 11	13–20 fl oz		
	<i>tebuconazole + azoxystrobin</i> Custodia Helmstar Plus	3 + 11	8.6-17.2 7.2 – 14.4	12 H/ 45 D	
	<i>tebuconazole + trifloxystrobin</i> Absolute	3 + 11	5-7.67 fl oz	12 H/ Do not apply after shuck split or within 30 D of harvest	
	<i>flutriafol + azoxystrobin</i> Topguard EQ	3 + 11	5.0 – 8.0 fl oz	12 H/ 45 D	
	<i>tetraconazole + triphenyltin hydroxide</i> Minerva Duo	3 + 30	16 oz	48 H/ 30 D	
	<i>thiophanate methyl</i> ³ (Topsin M) + TPTH or + Elast	1 + 30 or + U12	1 lb + half rate or + 25 fl oz	3 D/ Do not apply after shuck split	
	<i>triphenyltin hydroxide</i> (TPTH) ¹ + FRAC Group 3 fungicide	30 + 3	half rate ² + full rate	48 H/ 30 D	
Anthracnose	Anthracnose is a disease with a long latent period; symptom expression occurs many weeks after infection. Fungicides used for control of scab have been effective in suppressing anthracnose.				

PECAN DISEASE CONTROL

DISEASE	CHEMICAL & FORMULATION	MOA	RATE/ACRE	REI/PHI (Hours or Days)	COMMENTS
POST-POLLINATION APPLICATIONS: Every 10-21 Days From Nut Set To Shell Hardening					
Scab	<i>dodine</i> Elast 400F	U12	48 fl oz	48 H/ Do not apply after shuck split	<p>MOA Group 1: Risk for resistance is high. Use should be limited. When conditions are very favorable for scab, Topsin can be used in combination with either a full rate of TPTH or Elast. Limit the use to 1 or 2 applications per season.</p> <p>MOA Group 3: Resistance risk is moderate. For best results, tank mix <i>tebuconazole</i> with a surfactant. Do not add a surfactant if mixing with other fungicides. Increasing the rate of a Group 3 fungicide will be important if reduced sensitivity is known or suspected. Stand-alone use is not recommended where reduced sensitivity is known or suspected.</p> <p>MOA Group 11: Resistance risk is moderate. Do not make more than 2 sequential applications. If only using solo products, Group 11 fungicides should not be used in more than 1/3 of the total number of fungicide applications. If using Group 3 tank-mixed with other modes of action, they should not be used in more than 1/2 of the total number of fungicide applications.</p> <p>MOA Group 30: Resistance risk is low.</p> <p>MOA Group 33: Resistance risk is low. For best control apply in 100 gpa by ground. Do not apply in consecutive applications. Three to five applications are generally recommended. Check labels for potential limitations on maximum number of applications or amount of active ingredient allowed per season. Do not use when there is a phosphate deficiency.</p> <p>MOA Group U12: Resistance risk is low. Do not use on Moore, Van Deman, Barton, or Shawnee. Do not use a surfactant. Do not use with foliar zinc treatments.</p>
	<i>dodine</i> Elast 400F + FRAC Group 3 fungicide ³	U12 + 3	25 fl oz + full rate	48 H/ Do not apply after shuck split	
	<i>dodine</i> Elast 400F + TPTH	U12 + 30	25 fl oz + half rate ²	48 H/ Do not apply after shuck split	
	<i>phosphorous acid</i> Kphite 7LP Phostrol ProPhyt Viathon FungiPhite Reliant	33	highest label rate	4 H/ –	
	<i>phosphorous acid</i> + <i>tebuconazole</i> Viathon	33 + 3	2-2.5 pt	12 H/ 0 D	
	<i>propiconazole</i> + <i>azoxystrobin</i> Quilt Quilt Xcel	3 + 11 3 + 11	20-28 fl oz 20-21 fl oz	12 H/ Do not apply after shuck split or within 45 D of harvest	
	<i>tebuconazole</i> + <i>azoxystrobin</i> Custodia Helmstar Plus	3 + 11	8.6-17.2 7.2-14.4	12 H/ 45 D	
	<i>flutriafol</i> + <i>azoxystrobin</i> Topguard EQ	3 + 11	5.0 – 8.0 fl oz	12 H/ 45 D	
	<i>tebuconazole</i> ⁴ + <i>trifloxystrobin</i> Absolute	3 + 11	5-7.67 fl oz	12 H/ Do not apply after shuck split or within 30 D of harvest	
	<i>difenoconazole</i> + <i>azoxystrobin</i> Amistar Top	3 + 11	8-14 fl oz	12 H/ Do not apply after shuck split or within 30 D of harvest	
	<i>tetraconazole</i> + <i>azoxystrobin</i> Brixen	3 + 11	13–20 fl oz	12 H/ 45 D	
	tetraconazole + triphenyltin hydroxide Minerva Duo	3 + 30	16 oz	48 H/ 30 D	
	TPTH + FRAC Group 3 fungicide	30 + 3	half rate + full rate	48 H/ 30 D	

PECAN DISEASE CONTROL

DISEASE	CHEMICAL & FORMULATION	MOA	RATE/ACRE	REI/PHI (Hours or Days)	COMMENTS
POST-POLLINATION APPLICATIONS: Every 10-21 Days From Nut Set To Shell Hardening (continued)					
Scab (continued)	<i>triphenyltin hydroxide</i> (TPTH) ¹ Agri Tin Agri Tin Flowable Super Tin 80WP Super Tin 4L	30	7.5 oz 12 fl oz 7.5 oz 12 fl oz	48 H/ 30 D	See MOA info on previous page.
	<i>ziram</i> Ziram		6-8 lb	48 H/ 55 D	Ziram as a multi-site alternative in cases where resistance to other protectants is an issue.

Powdery Mildew	For powdery mildew, the scab fungicide program can be adjusted if needed. The FRAC Group 3 fungicides or mixes containing FRAC 3 fungicides are the best options. Combining sulfur (4-6 lb/A) with fungicides used for scab control is also an option. DO NOT mix sulfur with Elast.
Zonate Leaf Spot	For zonate leaf spot, the scab fungicide program can be adjusted if needed. The FRAC Group 3 fungicides or mixes containing FRAC 3 fungicides are the best options. Topsin M also provides suppression of Zonate leaf spot.
Anthracnose	Anthracnose is a disease with a long latent period; symptom expression occurs many weeks after infection. Fungicides used for control of scab have been effective in suppressing anthracnose, particularly FRAC Groups 3 and 11 and the phosphorous acid-based fungicides

¹ TPTH is available as Agri Tin, Agri Tin Flowable, Super Tin 80WP, and Super Tin 4L.

² Half rates are 3.75oz for Agri Tin and Super Tin 80WP; 6 fl oz for Agri Tin Flowable and Super Tin 4 L.

³ *Thiophanate methyl* is available as Topsin M 70WDG, Topsin M 70 WP, and Topsin M WSB, and Topsin M 4.5 FL (20 fl oz rate is equivalent to 1 lb of wettable powder). Topsin XTR is a premix of *thiophanate methyl* and *tebuconazole*.

⁴ For *tebuconazole*, use a minimum of 6 fl oz in tank mixes for nut scab control.

NOTE: In orchards where any nuts have any amount of scab by mid-June or in orchards where 10% or more of the nuts have any amount of scab by early July, the following measures should be taken:

- The interval between fungicide sprays should not exceed 14 days until shell hardening.
- On varieties with a summer growth flush, the spray interval should be closed so that no more than 10 days pass from the onset of the growth flush until a fungicide spray is made.
- If the 5-day forecast shows the probability for several days of rain, close the interval to have as much acreage as possible treated within 7 days of the storm.

PECAN DISEASE CONTROL

After Shell Hardening: Fungicide coverage for crop protection is necessary to shell hardening. Beginning in early August, monitor for shell hardening and adjust fungicide needs accordingly.

Foliar diseases: Maintaining leaf health past shell hardening is important. If leaf scab, zonate leaf spot, or another foliar disease is of concern, refer to the previous sections for fungicide options and recommendations. Pay attention to use limitations and fungicide resistance management guidelines. DO NOT use Topsin in consecutive applications for leaf disease control.

DISEASE	CHEMICAL & FORMULATION	MOA	RATE/ACRE	REI/PHI (Hours or Days)	COMMENTS
Phytophthora Shuck and Kernel Rot	A treatment is advised in orchards with a history of this disease (primarily Houston, Peach, and Macon counties) when wet weather and warm temperatures < 86° F occur between shell hardening and shuck split.				
	TPTH	30	full rate		
	<i>phosphorous acid</i> Fosphite Fungi-Phite KPhite Phiticide Phostrol Rampart Topaz	33	full rate	4 H/ –	The phosphite (phosphorous acid based) fungicides listed are EPA approved and considered to be very safe products. Check labels for potential limitations on maximum number of applications or amount of active ingredient allowed per season.
	MOA Group 11 fungicides	11	full rate		
	<i>copper hydroxide</i> Kocide 3000 Kocide 2000	M1	0.75-1.75 lb 1.5-3 lb	48 H/ –	Use higher rates when disease pressure is high and large, mature trees.

Restrictions and Fungicide Resistance Management Recommendations

- Follow label instructions for proper use of all fungicide products, including safe handling, tank mixing, application method, and resistance management.
- DO NOT apply more than 32 fl oz of *propiconazole*/A/season.
- DO NOT apply more than 32 fl oz of *tebuconazole*/A/season.
- DO NOT apply more than 1.5 qt of *fenbuconazole*/A/season.
- DO NOT use more than 45 oz of Agri Tin or Super Tin 80 WP or 72 fl oz of Agri Tin Flowable or Super Tin 4 L/A/season.
- DO NOT apply more than 1.6 lb (25.6 oz) of *kresoxim methyl*/A/season.
- DO NOT use Elast full season.
- If using a Group 3 fungicide alone pre-pollination, DO NOT use mixes containing a Group 3 fungicide postpollination.
- DO NOT make more than 2 sequential and 3 total applications of Group 11 fungicides.
- DO NOT apply more than 3 lb of *thiophanate methyl* (2.1 lb ai)/A/season.

COMMERCIAL PECAN WEED CONTROL

Wayne Mitchem, Extension Associate – Weed Science

Timothy Grey, Research Weed Scientist

HERBICIDE	MOA	BROADCAST RATE/ACRE		REI/PHI (Hours or Days)	REMARKS AND PRECAUTIONS
		AMOUNT OF FORMULATION	LBS ACTIVE INGREDIENT		
PRE-EMERGENCE					
<i>oryzalin</i> Surflan 4AS Oryzalin 4AS	3	2-6 qt	2-6		Use on non-bearing and bearing trees for control of annual grasses and small seeded broadleaf weeds. Use low rate for short-term control (2-4 months); high rate for long-term control (8-12 months). DO NOT apply to newly transplanted trees until soil has settled and no cracks are present. Apply before annual weeds emerge in the spring or add <i>paraquat</i> or <i>glyphosate</i> for control of emerged weeds. Sequential applications may be used so long as total use rate does not exceed 12 qt/A/year and there are 2.5 months between applications.
<i>diuron</i> Karmex XP or Diuron 80DF Direx or Diuron 4L other brands	7	2-4 lb 1.6-3.2 qt	1.6-3.2		Use for control of annual broadleaf weeds and some annual grasses only under trees established in the orchard at least 3 years. Apply in spring before annual weeds emerge; if weeds are present, include surfactant to improve contact activity. Make a single band or broadcast application as a directed spray. Use low rate on sandy loam soils. DO NOT use on sand, loamy sand, gravelly soils, or on exposed subsoils. DO NOT use on soils with less than 0.5% organic matter. DO NOT graze treated areas. Add <i>paraquat</i> , <i>glufosinate</i> , or <i>glyphosate</i> for enhanced control of emerged weeds.
<i>simazine</i> Princep, Simazine 90DF Princep, Simazine 4F	5	2.2-4.4 lb 2-4 qt	2-4		Use for control of annual broadleaf weeds and some annual grasses only under trees established for at least 2 years. Provides good control of annual ryegrass. Use low rates on sandy soils. DO NOT apply to gravelly, sand, or loamy sand soils. DO NOT apply when nuts are on the ground. DO NOT graze treated areas. Add <i>paraquat</i> , <i>glufosinate</i> , or <i>glyphosate</i> for control of emerged weeds.
<i>oryzalin</i> Surflan 4AS Oryzalin 4AS + <i>simazine</i> Princep, Simazine 80W 90DG 4L		2-4 qt + 2.5-5 lb 2.2-4.4 lb 2-4 qt	2-4 + 2-4		Use for broad spectrum annual grass and broadleaf weed control. Provides good control of annual ryegrass. <i>Paraquat</i> , <i>glufosinate</i> , or <i>glyphosate</i> may be used with this tank mix to enhance control of emerged weeds. See remarks and precautions for each product.
<i>norflurazon</i> Solicam 80DF + <i>diuron</i> Karmex 80DF Direx 4L		2.5-5 lb + 2-3.8 lb 1.6-3 qt	2-4 + 1.6-3		Use for broad spectrum annual grass and broad leaf weed control only under trees established in the orchard for at least 3 years. Apply in the spring before annual weeds emerge. See remarks and precautions for each product.
<i>pendimethalin</i> Prowl H ₂ O 4EC Prowl or Pendimethalin 3.3EC	3	2-6 qt 2.4-7.3 qt	2-6		Control of annual grasses and broadleaf weeds such as pigweed. Most effective when adequate rainfall or irrigation is received within 7 days after application. DO NOT apply to newly transplanted trees until ground has settled around roots. Sequential applications may be used as long as total use rate does not exceed 6 qt/A and there are 30 days between applications. Prowl H ₂ O has a 60 day PHI for pecans; however, other <i>pendimethalin</i> formulations can only be used in non-bearing pecans.

COMMERCIAL PECAN WEED CONTROL

HERBICIDE	MOA	BROADCAST RATE/ACRE		REI/PHI (Hours or Days)	REMARKS AND PRECAUTIONS
		AMOUNT OF FORMULATION	LBS ACTIVE INGREDIENT		
PRE-EMERGENCE					
<i>norflurazon</i> Solicam 80DF	12	2.5-5 lb	2-4		Use for control of annual grasses, broadleaf weeds, and suppression of some perennials under bearing, non-bearing, or newly set trees. Apply to newly planted trees only after soil has settled around roots, at least 6 months after planting. Avoid contact with roots. Apply in the fall or early spring–fall applications control a broader weed spectrum than spring applications. DO NOT apply when nuts are on the ground at harvest. Use low rate on coarse-textured soils, higher rates on fine-textured soils. Make only 1 application per year. DO NOT graze treated areas. May tank mix with <i>simazine</i> or <i>diuron</i> for broader spectrum weed control. Add <i>paraquat</i> , <i>glufosinate</i> , or <i>glyphosate</i> for control of emerged weeds. DO NOT apply within 60 days of harvest. Sequential applications can be used so long as total use rate does not exceed maximum use rate for soil texture and crop.
<i>rimsulfuron</i> Matrix 25WG Solida 25WG Previn 25WG	2	4 oz	0.063		Provide pre- and post- control of broadleaf and annual grass weeds (see label for weed control POST). For broad spectrum residual control tank mix with <i>diuron</i> , <i>oryzalin</i> , or Prowl H ₂ O. Use in orchards established at least 1 year. <i>Rimsulfuron</i> has a 14-day PHI for pecan. Sequential applications may be used so long as there are 30 days between applications and total use rate does not exceed 4 oz/A broadcast basis.
<i>flumioxazin</i> Chateau 51WDG Tuscany 51 WDG Tuscany 4 SC	14	6-12 oz	0.19-0.38		DO NOT apply more than 6 oz/A/application to soils having a sand and/or gravel content > 80%. Trees established less than 1 year must be shielded with a grow tube or waxed container. DO NOT apply second application within 30 days of initial application. Applications after bud break can only be made with shielded application equipment. Once trees break dormancy apply with <i>paraquat</i> or <i>glufosinate</i> for non-selective post-emergence control. Must use shielded application equipment if using in non-dormant pecan trees. <i>Flumioxazin</i> has a 60-day PHI for pecans.
<i>indaziflam</i> Alion 1.67SE	29	3.5-6.5 oz	0.045-0.085		Use in orchards established 3 years or longer. Sequential applications may be used as long as there are 90 days between applications and total use rate does not exceed 10.3 oz/A/year. Use rate cannot exceed 3.5 fl oz/A/application on soils having less than 1% organic matter. On soils with an organic matter content from 1-3%, no more than 5 fl oz/A can be applied in a single application and the total use rate for the year cannot exceed 8.5 fl oz/A. In order to apply more than 5 fl oz/A in a single application soil organic matter must be > 3%. Alion should be tank mixed with <i>glyphosate</i> , <i>glufosinate</i> , or <i>paraquat</i> for non-selective post-weed control. Alion has a 14-day PHI. Do not use on soils having a 20% or greater gravel content.
POST-EMERGENCE					
<i>2,4-D amine</i> Various generic formulations 3.8SL	4	2-3 pt	1-1.4		DO NOT apply more than twice a year or within 60 days of harvest. Trees must be at least 1 year old. DO NOT allow spray to drift onto or contact foliage, fruit, stems, or trunks of trees. DO NOT apply to bare ground. DO NOT apply on light, sandy soils. Past research has shown concerns of injury when applying <i>2,4-D</i> on sandy soils, immediately before a large rain and during early bud or leaf break. Extreme caution must be taken to avoid off target movement of <i>2,4-D</i> . Certain crops, like cotton and vegetables, can be severely injured by <i>2,4-D</i> drift. Some formulations may limit use rate 2 pt/A. Sequential applications may be used as long as there are at least 30 days between applications. See product label for details.

COMMERCIAL PECAN WEED CONTROL

HERBICIDE	MOA	BROADCAST RATE/ACRE		REI/PHI (Hours or Days)	REMARKS AND PRECAUTIONS
		AMOUNT OF FORMULATION	LBS ACTIVE INGREDIENT		
POST-EMERGENCE (continued)					
<i>fluzifop</i> Fusilade DX 2EC 2 lb/gal	1	8-24 fl oz	0.125-0.38		Use for control of annual and perennial grasses under bearing or non-bearing trees. Sequential applications will be necessary for control of perennial grass weeds like bermudagrass and johnsongrass. Low spray volumes (10 GPA) generally improve control. Add crop oil concentrate (1 qt/A). Make application to johnsongrass: 12-18" tall; bermudagrass: 3" tall or with 4-8" runners; annual grasses: 2-8" tall. Does not control nutsedge(s). DO NOT apply when harvestable nuts are on the ground. DO NOT graze treated area. DO NOT apply within 30 days of harvest.
<i>sethoxydim</i> Poast 1.5EC 1.5 lb/gal	1	1-2.5 pt	0.3-0.5		Use for control of annual and perennial grasses. Sequential applications will be necessary for control of perennial grass weeds like bermudagrass and johnsongrass. Low spray volumes (10 GPA) generally improve control. Add crop oil concentrate (1 qt/A). Use low rate on annual grasses up to 6" tall; higher rates on larger annual grasses and perennial grasses. Does not control nutsedge. DO NOT harvest within 15 days of application.
<i>clethodim</i> Select 2.0EC	1	6-8 fl oz			Use for control of annual and perennial grasses in NON-BEARING trees that will not be harvested within 1 year of application. Use higher rates and sequential applications for perennial grasses. Add a non-ionic surfactant containing at least 80% ai at a rate of 1 qt/100 gal of spray solution (0.25% v/v). Make application to johnsongrass: 12-18" tall; bermudagrass: 3" tall or with 4-8" runners; annual grasses: 2-8" tall. Does not control nutsedge.
Select Max 1 lb/gal					
Intensity One 1 lb/gal		12-1 6 oz			
<i>halosulfuron</i> Sanda 75WDG	2	0.67-1.33 oz	0.032-0.063		For control of nutsedge, pigweed, radish, and cocklebur. Apply as directed spray under trees established for at least 1 year. Avoid contact of spray with trunk, stem, roots, or tree foliage. May apply up to 2 applications. DO NOT apply within 1 day of harvest. See label for rate restrictions related to soil texture. Tank mix with <i>glyphosate</i> for broad spectrum control
<i>paraquat</i> Firestorm 3SL Parazone Paraquat Concentrate 3 lb/gal	22	1.75-2.7 pt	0.65-1		Use for broad spectrum, contact control of emerged weeds. Apply as a directed spray in at least 20 gal of water with 1-2 pt surfactant/100 gal of spray mix or 1% crop oil concentrate (1 gal/100 gal spray mix). Apply when annual weeds are succulent and 1-6" tall. DO NOT allow spray drift to contact foliage or green bark of trees since severe damage may occur. DO NOT allow animals to graze on treated areas. May be tank mixed with certain pre-emergence herbicides for effective residual weed control. DO NOT apply when nuts are on the ground.
Gramoxone SL 2 lb/gal		2-4 pt			
<i>glufosinate</i> Cheetah Reckon 280 Rely 280 Lifeline 2.34 lb/gal	10	48 fl oz	0.88-1.5		Use for broad spectrum control of emerged weeds and grasses, both annuals and perennials. Apply as a directed spray in high spray volumes on non-bearing and bearing trees. Possesses contact and limits systemic activity, but does well on wild brambles and perennial grasses. Does not have soil residual activity. DO NOT contact foliage or green bark. <i>Glufosinate</i> formulations are loaded with surfactant therefore NO additional nonionic surfactants or crop oil is needed. The addition of spray graded ammonium sulfate fertilizer at 8-10 lb/100 gal will enhance <i>glufosinate</i> activity.

COMMERCIAL PECAN WEED CONTROL

HERBICIDE	MOA	BROADCAST RATE/ACRE		REI/PHI (Hours or Days)	REMARKS AND PRECAUTIONS
		AMOUNT OF FORMULATION	LBS ACTIVE INGREDIENT		
POST-EMERGENCE (continued)					
<i>glyphosate acid</i> Numerous brands 4SL Roundup Weather Max 5.5SL	9	1-2 qt 11-46 fl oz	1-2		Use for broad spectrum control of emerged weeds, both annuals and perennials. Apply as a directed spray on bearing and non-bearing trees. DO NOT allow spray to contact foliage, suckers, or green bark of trees. Use low rate for control of annual weeds less than 12" tall. Refer to product label for rates to control specific perennial weeds. Repeat applications may be made. Some <i>glyphosate</i> formulations require the addition of an adjuvant. DO NOT allow <i>glyphosate</i> to contact bark or leaves. Try to avoid applications in late summer and fall. Trees are more sensitive to <i>glyphosate</i> during that time. Allow at least 3 days between last application and harvest.
<i>carfentrazone</i> Aim 2 lb/gal		0.5-2 oz	0.008-0.031		Apply alone or tank mix with other herbicides for post-emergence control of broadleaf weeds including pigweed, morningglory, lambsquarters and prickly lettuce. DO NOT allow Aim to contact desirable foliage, flowers, or fruit. DO NOT apply within 3 days of harvest. Trees less than 2 years old must be shielded from direct contact with Aim. Sequential applications may be used as long as total use rate does not exceed 7.9 oz/A/year and there are 14 days between applications. Best results obtained when applied to weeds in the 2-3 leaf stage. Apply in combination with a non-ionic surfactant (1 qt/100 gal of spray solution) or crop oil concentrate (1 gal/100 gal of spray solution).
ROW MIDDLE VEGETATION SUPPRESSION					
<i>glyphosate acid</i> Numerous brands 4SL Roundup Weather Max 5.5 S80 WDGL		2-16 fl oz 1.3-5.85	0.06-0.5 0.06-0.25		Use for vegetative suppression in row middles. Apply 1-2 weeks after full green-up of bahiagrass or bermudagrass, or after grass has been mowed to a uniform height of 3-4". Rates should vary depending on vigor of vegetative growth and canopy of the grove, with the higher rates for more vigorous grass stands where less shade occurs. Low spray volumes (10 GPA) improve control. See respective labels for surfactant requirements. Sequential applications can be made to maintain growth suppression and prepare the orchard floor for mechanical harvest. Allow a minimum of 21 days between the last application and harvest.

FOLIAR NICKEL SPRAYS FOR BEARING AND NON-BEARING PECAN TREES

TIME OF APPLICATION	MATERIAL	AMOUNT PER 100 GALLONS	INSTRUCTIONS AND REMARKS
Make application 10-24 days after bud break. Followed by a second application in mid-July.	Nickel Lignosulfonate (Nickel Plus) (6% Ni)	1-1.5 pt (April) 1.5-2 pt (July)	Can be tank-mixed with all fungicides, insecticides, nutrients, etc., including zinc. It is not necessary to add urea, which is already present.
For trees with visible mouse ear symptoms, or for newly transplanted trees, especially on sandy sites, or in orchards with a history of high zinc use, make a third application in early October before leaf fall.			Symptoms will be corrected 14-21 days after spring application, therefore all fall application ensures adequate levels of nickel in the plant tissue at bud- break. Research suggests that the lignosulfonate solution poses a lower risk for orchard workers and environmental safety than the metallic salt solution.
Make 1 st application at parachute stage and 2nd application 6 weeks later.	Nickel Sulfate (10% Ni)	1 pt	

All foliar micro-nutrient applications should be made only on an “as-needed” basis as determined by leaf tissue analysis and/or visual symptoms.

FOLIAR BORON APPLICATION FOR BEARING PECAN TREES

TIME OF APPLICATION	MATERIAL	AMOUNT PER 100 GALLONS	INSTRUCTIONS AND REMARKS
Begin Boron applications with 2nd Prepollination spray. Make 3 applications.	Solubur (20.5%)	1/16 lb of actual Boron	Do not concentrate. Do not apply more than 1 lb. of total Boron per season. When mixing Boron with imidacloprid, check pH of the solution and add a n acidifying agent when necessary to bring pH below 7.5.
	Boron Plus (10% B)	1 pt	
	Top Side Liquid Boron (6%)	1 pt	

NOTES

FORMULATION¹ ABBREVIATIONS

a.i. = active ingredient	EC = emulsifiable concentrate	S = sprayable powder
AC = aqueous concentrate	EL = emulsifiable liquid	SC = spray concentrate
AS = aqueous suspension	F = flowable	SP = soluble powder
DF = dry flowable	FC = flowable concentrate	W = wettable powder
DG = dispersible granules	G = granules	WDG = water dispersible granular
B = bait	L = liquid	WDL = water dispersible liquid
D = dust	LC = liquid concentrate	WM = water miscible
E = emulsifiable	M = microencapsulated	WP = wettable powder

¹ Numbers preceding abbreviations for liquid formulations equal pounds of active ingredient per gallon (e.g., 4EC = 4 lb/gal emulsifiable concentrate); numbers preceding abbreviations for solid formulations equal percent active ingredient by weight (e.g., 50WP = 50 percent wettable powder).

METHOD OR TIME OF APPLICATION ABBREVIATIONS

CR = cracking stage	PEI = pre-emergence incorporated	PRE = pre-emergence
LV = low volume	PO = post-emergence	PT = post-transplant
NS = nonselective	POT = post-emergence over-the-top	RCS = recirculating sprayer
PDS = post-emergence directed spray	PP = pre-plant	ULV = ultra low volume ²
PE = pre-emergence on surface	PPI = pre-plant soil incorporated	WICK = rope wick applicator

² Ultra low volume refers to a total spray volume of one-half gallon or less per acre.

RATE CALCULATIONS AND SPRAYER CALIBRATIONS

See the 2016 Georgia Pest Management Handbook or the “Pesticide Safety & Other Pesticide Information” section online at www.ent.uga.edu/pest-management/
--

MEASURES AND EQUIVALENTS

tsp	=	teaspoon		1 teaspoon	=	4.9 milliliters
Tbs.	=	tablespoon	1 Tbs	=	3 teaspoons	= 14.8 milliliters
fl oz.	=	fluid ounce	1 fl oz	=	2 tablespoons	= 29.6 milliliters
c	=	cup	1 c	=	8 fluid ounces	= 236.6 milliliters
pt	=	pint(s) (1.04 lb of water)	1 pt	=	2 cups	= 473.2 milliliters
pt/100	=	pint(s) per 100 gal	1 pt/100	=	1 teaspoon per gal	
qt	=	quart(s) (2.09 lb of water)	1 qt	=	2 pints	= 946.4 milliliters
gal	=	gallon(s) (8.35 lb of water)	1 gal	=	4 quarts	= 3.7854 liters
oz	=	ounce		1 ounce	=	28.35 grams
lb	=	pound	1 lb	=	16 ounces	= 453.59 grams
in	=	inch	1 in	=	1000 mils	= 2.54 centimeters (25,400 microns)
ft	=	feet	1 ft	=	12 inches	= 30.48 centimeters
yd	=	yard	1 yd	=	3 feet	= 91.44 centimeters
mi	=	mile	1 mi	=	5,280 feet	= 1,609 meters (16.09 kilometers)
sq in	=	square inch		1 square inch	=	6.45 square centimeters
sq ft	=	square feet	1 sq. ft	=	144 square inches	= 929.03 square centimeters
A	=	acre	1 A	=	43,560 square feet	= 0.4047 hectare
cu in	=	cubic inch		1 cubic inch	=	16.387 cubic centimeters
cu ft	=	cubic feet	1 cu ft	=	1,728 cubic inches	= 0.0283 cubic meter
cu yd	=	cubic yard	1 cu.yd	=	27 cubic feet	= 0.7646 cubic meter
ppm	=	parts per million	1 ppm	=	1,000 parts per billion	= 1 milligram/kilogram ³
psi	=	pounds per square inch		1 psi	=	70.3 gram-force per square centimeter

³ 1 milligram/kilogram or 1 p.p.m. is equal to 1 milligram/liter of water.

extension.uga.edu

Bulletin 841

Revised January 2019

Published by the University of Georgia in cooperation with Fort Valley State University, the U.S. Department of Agriculture, and counties of the state. For more information, contact your local UGA Cooperative Extension office.
The University of Georgia College of Agricultural and Environmental Sciences (working cooperatively with Fort Valley State University, the U.S. Department of Agriculture, and the counties of Georgia) offers its educational programs, assistance, and materials to all people without regard to race, color, religion, sex, national origin, disability, gender identity, sexual orientation or protected veteran status and is an Equal Opportunity, Affirmative Action organization.